



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ORBIT AND MASS OF THE VARIABLE STAR *ALGOL* (β *Persei*).

On the 28th of November a very important discovery was communicated to the Academy of Sciences of Berlin by Professor H. C. VOGEL, Director, and Dr. SCHEINER, Astronomer of the Astrophysikalisches Observatorium of Potsdam. I condense from the *Sitzungsberichte* of the Academy, 1889, (page 1045), the following :—

“Three photographic negatives of the spectrum of *Algol* taken during the winter of 1888-9 showed that before a minimum *Algol* was moving away from the sun, and after a minimum it was moving towards it. Three new exposures of November, 1889, confirm this result. The observations taken together afford a very strong support to the theory that the cause of the variations in the light of *Algol* is to be found in the eclipses of this star by a dark (invisible) satellite revolving about it. The phenomena can be explained by assuming the following particulars of the dimensions of the two bodies :—

- “ Diameter of *Algol* = 230,000 geographical miles.
- “ Diameter of the invisible satellite = 180,000 “ “
- “ Distance between their centres . = 700,000 “ “
- “ Satellite’s velocity in orbit . . = 12.0 “ “
- “ Mass of *Algol* = $\frac{4}{9}$ of the Sun’s mass.
- “ Mass of the satellite = $\frac{2}{9}$ “ “ “
- “ Motion of both bodies in the line of sight (toward the Sun) 0.5 geographical miles.”

E. S. H.

PROGRAMME FOR MERIDIAN OBSERVATIONS OF STARS.

When the REPSOLD meridian circle was ordered, it was designed to use it in determination of the absolute positions of the fundamental stars. Experiments during 1888 and 1889 showed Professor SCHAEFERLE (in charge of the circle) and myself that observations during the daytime were not at all comparable in precision with night observations. It is very rare to find the images in the daytime steady enough to deserve weight 2 (5 = perfectly steady, 1 extremely unsteady), while there are very many nights of weight 4 and weight 5 during the fair season (May to November). The Sun, *Mercury* and *Venus* were often observed at transit during 1889, and the images were always found to be of weight 1, or less. For this reason we reluctantly decided to abandon the plan of referring star places in Right Ascension to the Sun, either directly or through *Mercury* and *Venus*.

All our R. A. observations must then be differential.